

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425

1426

1427

1428

1429

1430

1431

1432

1433

1434

1435

1436

1437

1438

1439

1440

1441

1442

1443

1444

1445

1446

1447

1448

1449

1450

1451

1452

1453

1454

1455

1456

1457

1458

1459

1460

1461

1462

1463

1464

1465

1466

1467

1468

1469

1470

1471

1472

1473

1474

1475

1476

1477

1478

1479

1480

1481

1482

1483

1484

1485

1486

1487

1488

1489

1490

1491

1492

1493

1494

1495

1496

ROBERT F. FOX LIBRARY
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93940

NAVAL POSTGRADUATE SCHOOL

Monterey, California



THESIS

FUZZY PREFERENCE METHOD FOR DEFINING
GROUP PREFERENCES

by

Robert David Clarke

March 1980

Thesis Advisor:

Donald R. Barr

Approved for public release, distribution unlimited.

T191986

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Fuzzy Preference Method for Defining Group Preferences		5. TYPE OF REPORT & PERIOD COVERED Master of Science March 1980
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Robert David Clarke		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, California 93940		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Postgraduate School Monterey, California 93940		12. REPORT DATE
		13. NUMBER OF PAGES 45
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Naval Postgraduate School Monterey, California 93940		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) fuzzy sets, decision making, group preference ordering		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This paper presents a 'fuzzy preference' method of defining 'group preference' among alternatives based on the individuals' preferences. Group strengths of preference are also defined on the basis of individual strengths. An example and the results of four experiments are presented for the fuzzy preference method. Brief discussions are given of definitions and extensions in fuzzy set theory.		

Approved for public release; distribution unlimited

Fuzzy Preference Method for Defining Group Preferences

by

Robert David Clarke
Major, United States Marine Corps
B.S., United States Naval Academy, 1969

Submitted in partial fulfillment of the
requirements for the degree of

MASTER OF SCIENCE IN OPERATIONS RESEARCH

ABSTRACT

This paper presents a 'fuzzy preference' method of defining 'group preference' among alternatives based on the individuals' preferences. Group strengths of preference are also defined on the basis of individual strengths. An example and the results of four experiments are presented for the fuzzy preference method. Brief discussions are given of definitions and extensions in fuzzy set theory.

TABLE OF CONTENTS

	PAGE
I. INTRODUCTION	6
II. FUZZY SET THEORY	9
III. FUZZY PREFERENCE METHOD	16
IV. CONCLUSION	38
APPENDIX A: OUTLINE OF THE FUZZY PREFERENCE PROCEDURE.	41
LIST OF REFERENCES	43
INITIAL DISTRIBUTION LIST	45

LIST OF FIGURES

	PAGE
2.1 MEMBERSHIP FUNCTION FOR THE FUZZY SET B	10
2.2 MEMBERSHIP FUNCTIONS FOR THE FUZZY SETS A,B,C	12
2.3 α -LEVEL SETS OF A FUZZY SET D	13
2.4 MEMBERSHIP FUNCTIONS FOR FUZZY SETS A,B,C	15
3.1 INDIVIDUAL PREFERENCE MATRIX	21
3.2 INDIVIDUAL PREFERENCE MATRIX	21
3.3 COMPLETED INDIVIDUAL PREFERENCE MATRIX	22
3.4 AGGREGATION MATRIX	23
3.5 GROUP PREFERENCE MATRIX	24
3.6 DIAGRAM OF PREFERENCE ORDERING	25
3.7 DIAGRAM OF PREFERENCE ORDERING	25
3.8 DIAGRAMS OF PREFERENCE ORDERING	26
3.9 DIAGRAMS OF PREFERENCE ORDERING	27
3.10 AGGREGATION MATRIX WITH DOUBLE WEIGHTING	28
3.11 GROUP PREFERENCE MATRIX WITH DOUBLE WEIGHTING	29
3.12 DIAGRAMS OF PREFERENCE ORDERING WITH DOUBLE WEIGHTING	30
3.13 DIAGRAM OF PREFERENCE ORDERING WITH DOUBLE WEIGHTING	31
3.14 DIAGRAM OF PREFERENCE ORDERING WITH DOUBLE WEIGHTING	31
3.15 RESULTS OF FOOD EXPERIMENT	33
3.16 RESULTS OF SPORTS EXPERIMENT	34
3.17 RESULTS OF RETIREMENT EXPERIMENT	35
3.18 RESULTS OF MUSIC EXPERIMENT	36

I. INTRODUCTION

The goal of this writer is to present a simple, useable method of assisting decision-makers in solving a particular type of problem. The problem is how to use individual preferences to form the overall preferences of a 'group' of individuals. The reader having a modest background in mathematics should find this paper an interesting treatment of this type of problem. It is a problem often encountered in eliciting the group preference from a team of advisors. What are some of the difficulties faced by decision-makers in defining the preferences of a group of individuals over a set of alternatives?

The social, political, business, and military atmosphere today is clouded by a mood of complexity, ambiguity, and uncertainty. Increasingly more variables enter the arena; values change across the scope of time and personalities; the future continues to be quite unpredictable. Decision-makers are challenged to achieve more production with fewer expenditures. Commanders, managers, and analysts are frustrated by their inability to adequately capture the essence of their problems and to consistently render satisfactory solutions.

Zadeh [1] stated the 'principle of incompatibility' which gives us a sense of direction when we are confronted with the complexity of modern decision making.

"The closer one looks at a 'real world' problem, the fuzzier becomes its solution. Stated informally, the essence of this principle is that as the complexity of a system increases, our ability to make precise and yet significant statements about its behavior diminishes until a threshold is reached beyond which precision and significance (or relevance) become almost mutually exclusive characteristics."

According to Zadeh, fuzziness, vagueness, and imprecision are terms with shady connotations. Precision, logic, and clarity are terms perpetuated by the rigors of mathematics. "As we learn more about human cognition, we may well arrive at the realization that man's ability to manipulate fuzzy concepts is a major asset rather than a liability, and it is this ability, above all, that constitutes a key to the understanding of the profound difference between human intelligence, on one hand, and machine intelligence, on the other." [2] Therein lies the motivation for the language and logic of fuzzy sets.

Decision-making is an art and science of considerable scope. Extensive work has been done in many aspects of the theory and application of decision-making. The aim of this paper is to present just one of these aspects - preference ordering of alternatives for a group of individuals. Decision opportunities carry the goal of selecting from among alternatives the best choice or most favorable alternative. Implicit in the analytical process of making a choice is a preference ordering of alternatives. At least, a decision maker chooses one (or more) alternatives as best and the remainder as second

best. At most, the decision maker assigns a complete ordering of the alternatives according to his preferences. This process is inherently difficult for one person to do; for a group of people to arrive at a preferential consensus over a list of alternatives can be an arduous task. This paper presents a method for establishing a complete ordering of the alternatives for the group, based on the preferences of individual group members.

Section two of this paper presents some basic definitions of the theory for dealing with imprecision. Section three describes an application of fuzzy set theory, group preference ordering based on individual preferences. In section four there is a brief overview of literature and other applications of fuzzy set theory.

II. FUZZY SET THEORY

Fuzzy set theory was introduced by Zadeh for handling vague, inexact information in a mathematically rigorous way. In ordinary or 'crisp' set theory, each element of a particular set's universe of discourse either belongs or doesn't belong to the particular set. The universe of discourse is the set of all objects, defined by enumeration or rule, that will be considered in a given context. If, for instance, the universe of discourse is the set of positive integers, the set A of odd numbers less than ten is $A = \{1, 3, 5, 7, 9\}$. The set A is well defined, i.e., given any object it can be determined whether the object is in the set.

A fuzzy set taken over the same universe of discourse is the set B of odd numbers 'close to ten'. The characteristic function of a fuzzy set is allowed to take any value between 0 and 1 inclusive. Therefore, all elements in the universe of discourse 'belong' to a given fuzzy set, but with possibly different grades of membership. The membership function of the fuzzy set B is denoted by $\mu_B(x)$, where x varies over the universal set χ . Our set B might, for example, have the following membership function values:

$$\mu_B(9) = \mu_B(11) = 1.0$$

$$\mu_B(7) = \mu_B(13) = .75$$

$$\mu_B(5) = \mu_B(15) = .5$$

$$\mu_B(3) = \mu_B(17) = .25$$

$$\mu_B(1) = \mu_B(19) = \mu_B(21) = . . . = 0$$

The membership function for the fuzzy set B is represented graphically in Figure 2.1.




Figure 2.1 - MEMBERSHIP FUNCTION FOR THE FUZZY SET B

Fuzzy set theory is an extension of ordinary set theory. Basic definitions and relations to fuzzy set theory, such as union, intersection, and complementation, are extensions of the corresponding definitions in ordinary set theory.

The union of two fuzzy sets A and B on χ , denoted $A \cup B$, a fuzzy set, is defined by the membership function

$$\mu_{A \cup B}(x) = \max\{\mu_A(x); \mu_B(x)\}.$$

The intersection of two fuzzy sets A and B on χ , denoted $A \cap B$, is a fuzzy set defined by the membership function

$$\mu_{A \cap B}(x) = \min\{\mu_A(x); \mu_B(x)\}.$$

The complement of A, denoted \bar{A} , is a fuzzy set defined by

$$\mu_{\bar{A}}(x) = 1 - \mu_A(x).$$

Many algebraic properties from ordinary set theory also hold: commutativity, associativity, distributivity, and DeMorgan's theorems. Linguistic hedges such as 'very' and 'somewhat' operate on fuzzy membership functions to change their 'meanings'. Suppose, for example, the fuzzy set $A = \{\text{young}\}$ were defined by the membership function

$$\mu_A(x) = (1 + (.04x)^2)^{-1}$$

where the universe of discourse is ages of people, in a continuous sense. The membership function for the fuzzy set $B = \{\text{very young}\}$ might, for example, be given as the square of the former function.

$$\mu_B(x) = (1 + (.04x)^2)^{-2}.$$

On the other hand, suppose we define the membership function for fuzzy set $C = \{\text{somewhat young}\}$ as

$$\mu_C(x) = (1 + (.04x)^2)^{-1/2}.$$

The membership functions for these three contrived sets are graphically represented in Figure 2.2.




Figure 2.2 - MEMBERSHIP FUNCTIONS FOR THE FUZZY SETS A, B, C

The α -level set of a fuzzy set D on χ is defined as the ordinary (non-fuzzy) set $S_\alpha(D)$ for which the degree of membership in D exceeds or equals the level α :

$$S_\alpha(D) = \{x | \mu_D(x) \geq \alpha\}.$$

This is illustrated in Figure 2.3.




Figure 2.3 - α -LEVEL SETS OF A FUZZY SET D

A simple example follows that demonstrates an application of fuzzy sets in finding a fuzzy solution to a pair of competing objectives. In the example the objectives, expressed as fuzzy sets are

A: x should be much larger than 5

B: x should be near to 10

where:

$$\begin{aligned}\mu_A(x) &= 1 - (1 + (0.2(x - 5))^2)^{-1} && \text{for } x > 5 \\ &= 0 && \text{for } x \leq 5\end{aligned}$$

$$\mu_B(x) = (1 + (x - 10)^2)^{-1}$$

A question that might be asked is what values of x have positive membership in a fuzzy set that is much larger than 5 and near 10. What is the value for x that gives maximum membership value in this intersection? The intersection of the two fuzzy sets is the fuzzy set $C = \{\text{much larger than 5 and near 10}\}$, has the following membership function:

$$\mu_C(x) = \min\{\mu_A(x); \mu_B(x)\}$$

and is illustrated in Figure 2.4. All of the values of x greater than 5 have positive membership value. It can be seen on the membership function for C that all values of x greater than 5 have positive membership value. It can also be seen that x equal 10.85 has the maximum membership value. This example briefly shows one useful aspect of fuzzy set theory.




Figure 2.4 - MEMBERSHIP FUNCTIONS FOR FUZZY SETS A, B, C

This concludes a very brief overview of some definitions for fuzzy set theory. Fuzzy set theory is useful in defining group preferences based on individual preferences. This is discussed in section three. Extensions and a short literature review for fuzzy set theory are included in section four.

III. FUZZY PREFERENCE METHOD

Imprecision seems to be a characteristic of 'real world' problems. Fuzzy set theory appears to provide models useful in solving imprecise problems in a simple and useable manner. In this section we will outline a method of defining group preferences, based on the preferences of individuals within the group.

Group preference ordering is the collective ranking of alternatives, based on the preferences of individuals within the group. This paper describes only one of many ways to define group preferences. We assume that each group member can rank the alternatives according to his personal preferences. The problem at hand is to find an ordering of the alternatives consistent, in some sense, with the preferences of the 'group.' This problem occurs frequently in military, business, and social environments. Hierarchical structures lend themselves to staffs and committees which are tasked to present their preferences over the considered alternatives. What is a simple, useable method to define the preferences of a group?

Blin and Whinston [3] describe an appealing method for establishing and presenting the ranking of alternatives for a group, based on the rankings of individuals within the group. Generally, Blin and Whinston constructed a preference matrix A , composed of elements a_{ij} . Each element indicated the preference for the individual of alternative i over alternative j . In the Blin and Whinston model elements a_{ij}

were either 1 (alternative i is preferred over j) or 0 (alternative i is not preferred over j .) To establish preferences for the group, individual matrices were summed together to form an aggregation matrix B .

$$B = \sum_{k=1}^M A_k \quad \text{For } M = \text{number of individuals}$$

Blin and Whinston divided the aggregation matrix B by the number of individuals M to yield the group preference matrix C .

$$C = \frac{B}{M}$$

The elements C_{ij} of the group preference matrix C defined the group preference for the alternative i over alternative j . From the group preference matrix α -level sets were constructed to display 'agreement levels' for the group.

The fuzzy preference model presented in this paper is a variation of the Blin and Whinston method. The first difference in this model occurs in the construction of the individual group member preference matrix A . The elements of this matrix (a_{ij}) can assume values from 0 to 1.0 inclusive. This variation allows strengths of preference to be incorporated into the model. Strength of preference is a subjective measure of personal determination regarding the choice of one alternative over another. Strength of preference equal to one shows full resolve in the preference of one alternative

over another. Strength equal to zero indicates indifference in preference. The second difference in this paper is the weighting of individual preference matrices. Weighting allows the preferences and strengths of selected individual to have greater impact on the group preference ordering. In the weighted case the aggregation matrix is

$$B = \sum_{k=1}^M W_k A_k$$

where the W_k are weights for each individual. The group preference matrix becomes:

$$C = \frac{B}{\sum_{k=1}^M W_k} .$$

Aside from these two differences, the methodology and interpretation of our method remains the same as the Blin and Whinston model. An example and the results of four experiments are presented to demonstrate the simplicity of this method and suggest its usefulness. Four alternatives are treated in the example and experiments; a larger number could be readily managed. In small dimensional problems, the calculations required for this method are simple enough to be carried out by hand.

The first step of our fuzzy preference method is obtaining the preference ordering of alternatives by each individual group member. Two types of information should be

collected in this step. The first is a ranking of alternatives from most to least preferred. In the case where an alternative is not ranked, every element a_{ij} in the individual preference matrix carrying the subscript of the unranked alternative will be set to zero (row and column). The second type of information is the strength of preference between every two consecutive alternatives in the ranked list. If no members of a group recorded strengths of preference, the values should be set equal to one, as in the Blin and Whinston model. If a small number of group members did not record strengths of preference, values for their individual preference matrices may be set to one half or another arbitrary 'averaging' value. An alternate solution is reversion to the Blin and Whinston model with unity strengths for all individual preferences. An individual preference matrix with strengths of preference partially missing should be treated similar to the previous situation - averaging or reversion to unity strengths for all individual preferences.

Suppose, for example, that a person were ranking four alternate activities for Saturday afternoon. Alternatives might be:

- A. mow the lawn
- B. play tennis
- C. wash the car
- D. swim at the pool

An outline for ranking these alternatives could be:

> > >

The symbol '>' is interpreted 'is preferred to' and separates the positions for entering alternatives. For example, the following:

B > D > A > C

would be read; playing tennis is most preferred, followed by swimming, then mowing the lawn, and finally, washing the car. Continuing in the example, strengths of preference of an individual might be:

B > D > A > C
 .5 1.0 .2

Strength of preference can be any value between zero (no preference) and one (strong preference.) In our example, B and D are strongly preferred over A and C. Tennis is preferred over swimming, but with medium strength. Lawn mowing is preferred over washing the car, but only slightly.

The second step of fuzzy preference method is plotting the individual preferences from step one in a simple, square matrix format. The matrix is as large as the square of the number of alternatives being considered. In our example of activity preferences, the matrix has four rows and four columns.

	A	B	C	D
A				
B			*	
C				
D				

Figure 3.1 - INDIVIDUAL PREFERENCE MATRIX

A single element a_{ij} of this preference matrix A is interpreted as the individual's strength of preference of that row (representative alternative) over the column (representative alternative.) For example, the asterisk '*' position will contain the individual's strength of preference for B over C. Even before our example values are filled in we see that each diagonal value is zero, since an alternative has zero preference over itself. With our example values, construction of the preference matrix begins as follows:

	A	B	C	D
A	0			
B	1	0	1	.5
C			0	
D				0

Figure 3.2 - INDIVIDUAL PREFERENCE MATRIX

Entry of preferences of B over the other alternatives shows; B is preferred over A with strength 1.0, B is preferred over C with strength 1.0 and B is preferred over D with a strength .5 . Completing the example individual preference matrix yields Figure 3.3.

	A	B	C	D
A	0	0	.2	0
B	1	0	1	.5
C	0	0	0	0
D	1	0	1	0

Figure 3.3 - COMPLETED INDIVIDUAL PREFERENCE MATRIX

Strength of preference is a successive maximizing operation for each alternative. In our example the strength of preference for B over D is .5 while the strength of preference for B over A is $\max \{ .5, 1.0 \}$ or 1.0. The strength of preference for B over C is $\max \{ .5, 1.0, .2 \}$ or 1.0. This maximizing operation is pertinent to every successive strength of preference relationship. Alternative C is not preferred over any alternative. No alternative is preferred over B. At a glance, all preferences and strengths are available from the individual preference matrix.

Step three in our fuzzy preference method concerns aggregation of individual preference matrices. The aggregation

matrix B is constructed by summing the individual preference matrices.

$$B = \sum_{k=1}^M A_k$$

In the continuation of our example, suppose that six members of a group participated in the Saturday activity preference ordering. The following aggregation matrix B is contrived for the purpose of our example.

	A	B	C	D
A	0	.5	2.0	.3
B	5.5	0	5.3	2.0
C	1.5	.2	0	.7
D	5.6	2.1	5.0	0

Figure 3.4 - AGGREGATION MATRIX

Step four concerns the normalization of the aggregation matrix. In the case of equal weights among individual preference matrices the aggregation matrix B is divided by the number of group members M (in this example $M = 6$) to yield the group preference C.

$$C = \frac{B}{M}$$

The case of unequal weights is discussed later in this section.

Figure 3.5 is the group preference matrix for our example.

	A	B	C	D
A	0	.08	.33	.05
B	.92	0	.88	.33
C	.25	.03	0	.12
D	.93	.35	.83	0

Figure 3.5 - GROUP PREFERENCE MATRIX

Figure 3.5 displays information regarding preferences and strengths of preference as defined for the 'group.' Each element c_{ij} of the group preference matrix yields a sense of membership for alternative preference comparisons.

Step five of our method is to determine α -level sets for the group preference matrix C. Recalling from section two, an α -level set is the set of all elements having membership at least as large as the value of α . In the example the α -level set at $\alpha = 1.0$ is empty, as can be seen from Figure 3.5. At $\alpha = .93$ there is one preference, namely D is preferred over A (with strength of preference = .93.) The α -level set for $\alpha = .93$ has one element, D preferred over A. This set is denoted as follows:

$$R_{.93} = \{(D,A)\}.$$

As you recall, the alternatives are:

- A. mow the lawn
- B. play tennis
- C. wash the car
- D. swim at the pool

Figure 3.6 should aid in understanding the preference ordering.




Figure 3.6 - DIAGRAM OF PREFERENCE ORDERING

At $\alpha = .92$ a second relationship occurs and the α -level set at $\alpha = .92$ has two members - (D,A) and (B,A).

$$R_{.92} = \{(D,A) , (B,A)\}$$





Figure 3.7 - DIAGRAM OF PREFERENCE ORDERING

Figure 3.7 indicates that alternative A is strongly not preferred. So far, we have not defined a complete ordering for every alternative for the group.

Subsequent α -level sets are constructed in like manner.

$$R_{.88} = \{(D,A), (B,A), (B,C)\}$$



$$R_{.83} = \{(D,A), (B,A), (B,C), (D,C)\}$$





Figure 3.8 - DIAGRAMS OF PREFERENCE ORDERING

At this point we see that the group prefers B and D over A and C. There is yet to appear, in our α -level set constructions, a relationship of A with respect to C and B with respect to D.

$$R_{.35} = \{(D,A), (B,A), (B,C), (D,C), (D,B)\}$$



$$R_{.33} = \{(D,A), (B,A), (B,C), (D,C), (D,B), (A,C)\}$$




Figure 3.9 - DIAGRAMS OF PREFERENCE ORDERING

A complete ordering of the four alternatives is revealed in $R_{.33}$. For the example the fuzzy preference of the group is

$$\begin{array}{cccc} D & > & B & > & A & > & C \\ .35 & & .92 & & .33 & & \end{array}$$

The values in the blocks are the strengths that accompany the fuzzy preferences. Strong preferences were defined for D and B over A and C, but weak preferences for D over B and A over C.

A variation to this method allows the decision-maker to assign weighting factors to those group members whose opinions are more valued by him for a particular set of alternatives. In our example, suppose that one member's preference matrix is weighted by a factor of two. In this sense, the particular member's preferences are twice as valuable to the

decision-maker as any other member. Recalling from before, the aggregation matrix for weighting is

$$B = \sum_{k=1}^M W_k A_k.$$

Figure 3.10 is a contrived aggregation matrix for the Saturday activity example, with double weighting for one group member.

	A	B	C	D
A	0	.5	2.4	.3
B	6.4	0	6.3	2.3
C	1.5	.2	0	.7
D	6.0	2.1	5.4	0

Figure 3.10 - AGGREGATION MATRIX WITH DOUBLE WEIGHTING


Step four normalizes the aggregation matrix B by dividing by the sum of the weights ($\sum_{k=1}^M W_k = 7$, in our double-weighted example.) The weighting factor gave the result as if a seventh group member were casting a preference vote equal to the 'more valuable' member preference. The group preference matrix with weighting is shown in Figure 3.11.

	A	B	C	D
A	0	.07	.34	.04
B	.91	.0	.90	.33
C	.21	.03	0	.10
D	.86	.30	.77	0



Figure 3.11 - GROUP PREFERENCE MATRIX WITH DOUBLE WEIGHTING

The new α -level sets are similar to the unweighted example with an important exception at $\alpha = .33$. Figure 3.12 shows the construction of the weighted α -level sets.



$$R_{.91} = \{(B, A)\}$$



$$R_{.90} = \{(B, A), (B, C)\}$$



$$R_{.86} = \{(B, A), (B, C), (D, A)\}$$



$$R_{.77} = \{(B, A), (B, C), (D, A), (D, C)\}$$




Figure 3.12 - DIAGRAMS OF PREFERENCE ORDERING WITH DOUBLE WEIGHTING

At this point we are quite close to the unweighted case where $\alpha = .83$.

$$R_{.34} = \{(B,A), (B,C), (D,A), (D,C), (A,C)\}$$




Figure 3.13 - DIAGRAM OF PREFERENCE ORDERING WITH DOUBLE WEIGHTING

At $\alpha = .33$ the double weighting of a member (who preferred B over D) causes a reversal in the group's fuzzy preference between B and D.

$$R_{.33} = \{(B,A), (B,C), (D,A), (D,C), (A,C), (B,D)\}$$




Figure 3.14 - DIAGRAM OF PREFERENCE ORDERING WITH DOUBLE WEIGHTING

$R_{.33}$ defines a complete ordering of the four alternatives.

$$\begin{array}{ccccccc} B & > & D & > & A & > & C \\ .33 & & .86 & & .34 & & \end{array}$$

The preference ordering of B and D has changed. The strengths of fuzzy preferences are shown in Figure 3.11. The preference

of B over D has strength of .33, while the preference of D over B has strength of .30 . B is only weakly preferred over D, but B and D are strongly preferred over A and C .

An outline of the fuzzy preference method is contained in Appendix A.

Experiments were conducted to test the simplicity and usefulness of the fuzzy preference method. Sets of alternatives were chosen so as to be familiar to the participants and to generate heterogeneous results. The fuzzy preference method was exercised in the unweighted mode by twelve volunteer graduate students at the Naval Postgraduate School. The results of four experiments, each with four alternatives, are shown in Figures 3.15 - 3.18. Each figure gives the alternatives, group preference matrix, α -level set at which complete ordering of alternatives is achieved, and the fuzzy preferences of alternatives.

The first experiment was a choice of foods:


Alternatives

- A. Roast Leg of Lamb
- B. Prime Rib of Beef
- C. King Crab
- D. Lobster

Group Preference Matrix

	A	B	C	D
A	0	.08	.18	.13
B	.64	0	.51	.43
C	.57	.26	0	.08
D	.66	.32	.46	0

$$R_{.43} = \{(B,D), (B,C), (B,A), (D,C), (D,A), (C,A)\}$$



B > D > C > A
 .43 .46 .57

Figure 3.15 - RESULTS OF FOOD EXPERIMENT

The second experiment was a preference ordering of participant sports:


Alternatives

- A. Water Skiing
- B. Tennis
- C. Bowling
- D. Roller Skating

Group Preference Matrix

	A	B	C	D
A	0	.28	.41	.54
B	.39	0	.68	.63
C	.32	.18	0	.53
D	.19	.22	.06	0

$$R_{.39} = \{(B,A), (B,C), (B,D), (A,C), (A,D), (C,D)\}$$



B >

A >

C >

D

.39

.41

.53

Figure 3.16 - RESULTS OF SPORTS EXPERIMENT

The third experiment was preference ordering of retirement location.


Alternatives

- A. Northeast
- B. Northwest
- C. Southeast
- D. Southwest

Group Preference Matrix

	A	B	C	D
A	0	.04	.15	.12
B	.73	0	.40	.39
C	.50	.29	0	.38
D	.64	.20	.30	0

$$R_{.38} = \{(B, C), (B, D), (B, A), (C, D), (C, A), (D, A)\}$$



$B > C > D > A$
 .40 .38 .64

Figure 3.17 - RESULTS OF RETIREMENT EXPERIMENT

The fourth experiment was preference ordering of music:

Alternatives

- A. Rock and Roll
- B. Disco
- C. Classical
- D. Jazz

Group Preference Matrix

	A	B	C	D
A	0	.25	.22	.35
B	.13	0	.12	.30
C	.48	.57	0	.69
D	.27	.34	.08	0

$R_{.34} = \{(C,A), (C,D), (C,B), (A,D), (A,B), (D,B)\}$




Figure 3.18 - RESULTS OF MUSIC EXPERIMENT

The results of the example problem and four experiments reveal the simplicity of the fuzzy preference method. Usefulness of this method is suggested by these applications. Defining the preference of a group can be a valuable aid to decision-makers. Strength of preference helps to show the decision-maker the magnitude of group support for each preference.

In the concluding section of this paper we briefly discuss extensions and criticism of this model and introduce literature for further study.

IV. CONCLUSION

In concluding this presentation we look backward and forward. First, did we answer our initial questions? We have found an intuitively appealing aid for decision-makers to elicit group preferences based on individual preferences over a set of alternatives. It is a simple, useable model that shows order and strength of preferences for a group of individuals.

Critically, the greatest possibility of difficulty occurs in group members making their preferences and strengths of preferences. This process is highly dependent on subjective judgement and exhibits the strong and weak characteristics of subjectivity. Favorably, subjectivity allows a person to bring to bear on the problem quantitative and qualitative information, experience, and intuition. Unfavorably, subjectivity invites personal prejudice and unrelated bias to affect the outcome. Also, the number of alternatives should be small enough to allow group members to ascertain their preferences over the entire set of alternatives.

With this model, decision-makers have the option of biasing the group preferences by weighting the preferences of selected members of his group. In a problem, for instance, oriented towards operations a commander might weight the preferences of the operations staff officer higher than those of his other group members.

Looking ahead, our method can be extended in many directions. Different weighting schemes would give decision-makers flexibility in determining group preferences. Another approach is an extension that would allow group members to assess their own level of expertise. With a high self-assessment, a member would weigh his preference higher, compared to fellow members. With a low self-assessment, a member weights in a way to detract from his preference contribution to the group aggregation matrix.

Considerable work has been done in fuzzy set theory. Zadeh pioneered the study of fuzzy sets with a paper in 1965 [4]. Since then he has published papers on linguistic variables, approximate reasoning, and fuzzy languages [5]. Bellman and Zadeh published a paper in 1970 that provided a basis for multi-criteria decision making in fuzzy sets [6]. A more recent presentation on multicriteria decision making was published by Blin [7]. Fuzzy preference functions have been investigated by Roy [8] and Bezdek [9]. Mathematical programming, considering decision problems with special structure, has been presented by Tanaka, Okuda, Asai [10], and Zimmermann [11]. Yager [12] has considered general decision making in fuzzy environments.

Fuzzy set theory has found application in many areas besides decision theory. Fuzzy methodologies in matching have been applied to criminal investigation [13], personnel management [14], and information processing [15]. Artificial

intelligence [16] and fuzzy control [17,18] are areas explored in fuzzy linguistics. Medical diagnosis using fuzzy set theory has been done by Esogbue [19] and Wechsler [20]. Applications of fuzzy set theory have also been made to psychological problems [21]. A bibliography written by Gaines and Kohout [22] is an excellent guide through fuzzy set theory and it's applications.

APPENDIX A

OUTLINE OF THE FUZZY PREFERENCE PROCEDURE

Step 1 Each group member records his preferences and strength of preferences over the alternatives.

> > >

Step 2 Plot each record of preference in a square matrix format.

	A	B	C	D
A	0			
B		0		
C			0	
D				0

Step 3 Combine individual preference matrices with weights in one aggregation matrix.

$$B = \sum_{k=1}^M w_k A_k$$

Step 4 Normalize the aggregation matrix to create the group preference matrix

$$C = \frac{B}{\sum w_k}$$

Step 5 Determine the α -level sets from the group preference matrix. Finally, indicate the complete ordering of the alternatives.

LIST OF REFERENCES

1. Zadeh, L. A., "Outline of a New Approach to the Analysis of Complex Systems and Decision Process," IEEE Transactions on Systems, Man, and Cybernetics, p. 28, January 1973.
2. Gupta, M. M., Saridis, G. N., and Gaines, B. R., eds., Fuzzy Automata and Decision Processes, North Holland, p. 3, 1977.
3. Blin, J. M. and Whinston, A. B., "Fuzzy Sets and Social Choice," Journal of Cybernetics, v. 3, p. 28-36, 1974.
4. Zadeh, L. A., "Fuzzy Sets," Information and Control, v. 8, p. 338-353, 1965.
5. Zadeh, L. A., "Concept of a Linguistic Variable and its Application to Approximate Reasoning," Information Science, v. 8, p. 199, 1975.
6. Bellman, R. and Zadeh, L. A., "Decision-Making in a Fuzzy Environment," Management Science, v. 17, p. 141-164, 1970.
7. Blin, J. M., "Fuzzy Sets in Multiple Criteria Decision-Making," TIMS Studies in the Management Sciences, v. 6, p. 129-116, 1977.
8. Roy, B., "How Outranking Relation Helps Multiple Criteria Decision-Making," Multiple Criteria Decision-Making, ed. by Cochrane, J. L. and Zeleny, M., University of South Carolina Press, p. 179-201, 1973.
9. Bezdek, J. C., Spillmann, B., and Spillmann, R., "A Fuzzy Relation Space for Group Decision Making," Fuzzy Sets and Systems, v. 1, n. 4, p. 255-268, 1978.
10. Tanaka, H., Okuda, T., and Asai, K., "On Fuzzy Mathematical Programming," Journal of Cybernetics, v. 3, p. 37-46, 1974.
11. Zimmermann, H. J., "Description and Optimization of Fuzzy Systems," International Journal of General Systems, v. 2, p. 209-215, 1976.
12. Yager, R. R. and Basson, D., "Decision Making with Fuzzy Sets," Decision Science, v. 6, p. 590-600, 1975.

13. Van Velthoven, G. D., Application of Fuzzy Set Theory to Criminal Investigation, Ph.D. Thesis, University of Louvain, Belgium, 1974.
14. Van Velthoven, G. D., "Fuzzy Models in Personnel Management," Procedures of Third International Congress of Cybernetics and Systems, Bucharest, 1975.
15. Zimmermann, H. J. and Gehring, H., "Fuzzy Information Profiles for Information Selection," Fourth INTERNET Congress AFCET, Paris, 1975.
16. Assilian, S., Artificial Intelligence in the Control of Real Dynamic Systems, Ph.D. Thesis, Queen Mary College, University of London, 1974.
17. Mamdani, E. H., "Advances with Linguistic Synthesis of Fuzzy Controllers," International Journal of Man-Machine Studies, v. 8, p. 669-678, 1976.
18. Kickert, W.J.M. and Van Nauta Lemke, H. R., "Application of Fuzzy Controller in a Warm Water Plant," Automatica, v. 12, p. 301-308, 1976.
19. Esogbue, A. O. and Elder, R. C., Fuzzy Sets and the Modelling of Physician Decision Processes, Parts I and II, Industrial and Systems Engineering Report Series No. J-77-6, Georgia Institute of Technology, February 1977.
20. Wechsler, "Applications of Fuzzy Logic to Medical Diagnosis," Procedures of 1975 International Symposium of Multiple-Valued Logic IEEE 75CH959-7C, May 1975.
21. Kochen, M., "Applications of Fuzzy Sets in Psychology, "Fuzzy Sets and their Applications to Cognitive and Decision Processes, ed. by Zadeh, L. A., Fu, K. S., Tanaka, K., and Shimura, M., p. 395-408, Academic Press, 1975.
22. Gaines, B. R. and Kohout, L. J., "The Fuzzy Decade: A Bibliography of Fuzzy Systems and Closely Related Topics," International Journal of Man-Machine Studies, v. 9, p. 1-68, 1977.

INITIAL DISTRIBUTION LIST

	<u>No. Copies</u>
1. Defense Technical Information Center Cameron Station Alexandria, Virginia 22314	2
2. Library, Code 0142 Naval Postgraduate School Monterey, California 93940	2
3. Department Chairman, Code 55 Department of Operations Research Naval Postgraduate School Monterey, California 93940	1
4. Professor D. R. Barr, Code 55Br (thesis advisor) Department of Operations Research Naval Postgraduate School Monterey, California 93940	1
5. Associate Professor G. F. Lindsay Code 55Ls (second reader) Department of Operations Research Monterey, California 93940	1
6. Maj. R. David Clarke, USMC (student) 9273 Bayberry Avenue Manassas, Virginia 22110	1

Thesis

C4975 Clarke


c.1

Fuzzy preference
method for defining
group preferences.

187080

thesC4975

Fuzzy preference method for defining gro



3 2768 001 02619 8

DUDLEY KNOX LIBRARY